

# Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## PROBLEMS AND SOLUTIONS.

EDITED BY B. F. FINKEL AND R. P. BAKER.

#### PROBLEMS FOR SOLUTION.

#### ALGEBRA.

When this issue was made up, no solution had been received for number 428.

## 430. Proposed by H. C. FEEMSTER, York College, Nebraska.

Solve the equations

$$\sum_{i=1}^{n} x_i - x_n = k + \frac{n^2 - 3n + 2}{2} d, \tag{1}$$

$$\sum_{i=1}^{n} x_i - x_{n-1} = k + \frac{n^2 - 3n + 4}{2} d, \tag{2}$$

$$\sum_{i=1}^{n} x_i - x_{n-2} = k + \frac{n^2 - 3n + 6}{2} d, \tag{3}$$

$$\sum_{i=1}^{n} x_i - x_1 = k + \frac{n^2 - n}{2} d.$$
 (n)

## 431. Proposed by ELMER SCHUYLER, Brooklyn, N. Y.

Form a magic square of 9 cells such that (the integers being all different) the products of the integers in the rows, columns, and diagonals shall be the same and the smallest product possible.

#### 432. Proposed by C. N. SCHMALL, New York City.

There are n straight lines in a plane, no two of which are parallel, and no three of which are concurrent. Their points of intersection being joined, show that the number of new lines drawn is  $\frac{1}{3}n(n-1)(n-2)(n-3)$ .

## GEOMETRY.

When this issue was made up, no solution had been received for number 459.

#### 460. Proposed by J. W. CLAWSON, Ursinus College, Pa.

ABC is a triangle, O and I the centers of the circum- and in-circles respectively, and I', I'', I''' the centers of the three escribed circles. If AO, BO, CO meet the circumcircle in P, P', P'' respectively, and PR, P'R', P''R'' are drawn parallel respectively to AI, BI, CI to meet BC, CA, AB respectively in R, R', R'', prove that: (1) PR, P'R', P''R'' are concurrent, say at I. (2) JO = OI. (3) JI', JI'', JI''' are perpendicular respectively to BC, CA, AB. (4) AR, BR', CR'' are concurrent.

### 461. Proposed by CLIFFORD N. MILLS, Brookings, South Dakota.

Prove by means of any inscribed triangle the following trigonometrical relations:  $\sin 2\beta = 2 \sin \beta \cos \beta$ .  $\cos 2\beta = \cos^2 \beta - \sin^2 \beta$ .  $\sin 3\beta = 3 \sin \beta - 4 \sin^3 \beta$ .  $\cos 3\beta = 4 \cos^3 \beta - 3 \cos \beta$ .

## 462. Proposed by DANIEL KRETH, Wellman, Iowa.

A conical glass, the diameter of the base of which is 5 inches and altitude 6 inches, is one-fifth full of water. If a sphere 4 inches in diameter is dropped into it, how much of the vertical axis of the glass is immersed?

## 463. Proposed by NATHAN ALTSHILLER, University of Washington, Seattle.

Through a given point, to draw a line that cuts off on the sides of a given angle two segments  $_{+}$ he sum of which has a given value.